



JONAS R. RUDZKI, MD

SPORTS MEDICINE ORTHOPAEDIC SURGEON,

SPECIALIZING IN SHOULDER & KNEE SURGERY

WASHINGTON ORTHOPAEDICS AND SPORTS MEDICINE

K STREET/DOWNTOWN

Phone: 202-833-1147

Fax: 202-296-2515

2021 K St NW, Suite 516

Washington, DC 20006

CHEVY CHASE

Phone: 301-657-1996

Fax: 301-951-6160

5454 Wisconsin Ave

Suite 1000

Chevy Chase, MD 20815

SIBLEY

Phone: 202-787-5601

Fax: 202-787-5606

5215 Loughboro Rd NW

Suite 200

Washington, DC 20016

POST-OPERATIVE PATIENT INSTRUCTIONS:

Coracoclavicular/Acromioclavicular Reconstruction

1. **Before surgery, please call Dr. Rudzki's MA to schedule your first postop appointment for 7-10 days post-op.** At the 1st post-op appointment, we will remove your dressings, remove your sutures, and discuss the plan for your therapy and recovery. If you have staples, they will be removed at your 2nd postop, and we will determine the timeframe for the 2nd post-op appointment your 1st post-op appointment.
2. If you develop a fever (temperature greater than 101.5 F), redness or drainage from the surgical incision site, please call our office to arrange for an evaluation.
3. You will receive two prescriptions for pain medication (typically morphine sulfate & Oxycodone).
 - These pain medications are all opioid narcotic derivatives, which have potential side effects that include constipation, nausea and sleepiness. Please take over the counter laxatives and stool softeners (ex. Peri-Colace, MiraLax, etc.), drink a lot of water, and eat dried apricots/ prunes 3x/day. These medications should not be taken on an empty stomach.
 - Please **do not take any Motrin, Ibuprofen, or Aleve (NSAIDs)** as these may impair your reconstructed ligament from healing.
4. You may slowly resume a regular diet as tolerated. We recommend you start slowly with clear liquids, diluted Gatorade, chicken/beef broth & crackers:
 - As soon as you get home, take small bites of food (1/4 turkey sandwich, diluted Gatorade, chicken soup, crackers) and 30 minutes later take your first pain pill (Oxycodone). Repeat this process 45-60 minutes later and take a 2nd pain pill (Oxycodone). Repeat again 45-60 minutes later and take a morphine sulfate. From this point forward, take Oxycodone 1-2 tabs every 4-6 hrs and morphine sulfate 1 tab every 12 hours. **It's important to stay ahead of pain, but do not take the pain pills if you are sedated. If you are 21 years or older, take 1 baby aspirin twice a day for DVT prophylaxis.**
5. Remove the dressings on the 2nd day after surgery. Apply Band-Aids loosely over the incisions and change them every 2-3 days as needed. Please do not use Bacitracin, Neosporin or any other creams under the bandages.
 - **Do not shower or get your incision wet until cleared to do so by Dr. Rudzki.**
6. Apply the Game Ready Ice as often as possible for the first 10-14 days; then at least 4-6 times each day for 30 minutes and after each time you do your exercises. The Game Ready Unit should be set to cool for 30 minutes & then go off for 30 minutes. Take a one hour break every 4-6 hrs. **Make sure incisions stay dry.**
7. Your sling should be worn at all times until your first post-operative visit to provide both comfort and support.
 - Come out of the sling 2 times a day for gentle elbow & wrist range-of-motion exercises with your arm supported at the side. Please work on wrist & finger range-of-motion exercises when in the sling.
 - It is helpful to sleep propped up with pillows and with a pillow supporting your elbow.
8. If you had a hamstring tendon harvest to augment your reconstruction, you will follow the same instructions for your knee incision. Remove the dressing on the 2nd morning after surgery. Ice your knee 4x/day for ~30 minutes. Elevate your leg whenever possible. You may bear weight on your leg without limitation as tolerated. Keep the incision clean & dry at all times.

Dr. Rudzki's MA can be reached at 202-787-5601 ext 652 or RudzkiMA@alignedortho.com



Website: www.jrrudzkimd.com