



JONAS R. RUDZKI, MD

SPORTS MEDICINE ORTHOPAEDIC SURGEON,
SPECIALIZING IN SHOULDER & KNEE SURGERY

WASHINGTON ORTHOPAEDICS AND SPORTS MEDICINE

Post-Operative Instructions: Knee Arthroscopy

Arthroscopic Partial Meniscectomy / Chondroplasty / Debridement

K STREET/DOWNTOWN

Phone: 202-833-1147

Fax: 202-296-2515

2021 K St NW, Suite 516

Washington, DC 20006

CHEVY CHASE

Phone: 301-657-1996

Fax: 301-951-6160

5454 Wisconsin Ave

Suite 1000

Chevy Chase, MD 20815

SIBLEY

Phone: 202-787-5601

Fax: 202-787-5606

5215 Loughboro Rd NW

Suite 200

Washington, DC 20016



1. **Before surgery, please call Dr. Rudzki's MA to schedule your first postop appointment for 7-10 days post-op.** At the 1st post-op appointment, we will remove your dressings, remove your sutures, and discuss the plan for your therapy and recovery. If you have staples, they will be removed at your 2nd postop, and we will determine the timeframe for the 2nd post-op appointment your 1st post-op appointment.
2. If you develop a fever (temp > 101.5°), redness or drainage from the surgical incision site, please call our office to arrange for a prompt evaluation.
3. You will receive two prescriptions for pain medication (typically morphine sulfate & Oxycodone).
 - These pain medications are all opioid narcotic derivatives, which have potential side effects that include constipation, nausea and sleepiness. Please take over the counter laxatives and stool softeners (ex. Peri-Colace, MiraLax, etc.), drink a lot of water, and eat dried apricots/ prunes 3x/day. These medications should not be taken on an empty stomach.
 - Please **do not take any Motrin, Ibuprofen, or Aleve (NSAIDs)** as these may impair your reconstructed ligament from healing.
4. You may slowly resume a regular diet as tolerated. We recommend you start slowly with clear liquids, diluted Gatorade, chicken/beef broth & crackers:
 - As soon as you get home, take small bites of food (1/4 turkey sandwich, diluted Gatorade, chicken soup, crackers) and 30 minutes later take your first pain pill (Oxycodone). Repeat this process 45-60 minutes later and take a 2nd pain pill (Oxycodone). Repeat again 45-60 minutes later and take a morphine sulfate. From this point forward, take Oxycodone 1-2 tabs every 4-6 hrs and morphine sulfate 1 tab every 12 hours. **It's important to stay ahead of pain, but do not take the pain pills if you are sedated.** If you are 21 years or older, take 1 baby aspirin twice a day for DVT prophylaxis.
5. Remove the dressing the day after surgery. Apply Band-Aids to the incisions(s). Please do not use Bacitracin, Neosporin or any other creams under the bandages.
 - **Do not shower or get your incision wet until cleared to do so by Dr. Rudzki.** Please do not bathe/soak the surgical area or go swimming in the pool/hot tub/ocean.
6. Elevation and protection of your surgical site are the most important considerations for initial post-op care. Keep your leg elevated with a 3-4 pillows under your calf & ankle, NOT under the knee.
7. You may use crutches as needed for stability but you are allowed to bear weight as tolerated on your leg. As you work on strengthening your leg and decreasing your swelling, you will be able to discontinue your use of the crutches.
8. You may experience some low back pain due to muscle spasm from the epidural anesthesia. If so, apply heating pad to area and take an analgesic if you have not already done so.

Dr. Rudzki's MA can be reached at 202-787-5601 ext 652 or RudzkiMA@alignedortho.com

Website: www.jrrudzkimd.com